

Breakthrough into the Sub-eV Neutrino Mass Range: Status of the HEIDELBERG-MOSCOW Double Beta Decay Experiment with enriched ^{76}Ge

A.Balysh^b, M.Beck^a, S.T.Belyaev^{*,b}, J.Bockholt^a, A.Demehin^b, M.Eskef^a, D.Glatting^a,
A.Gurov^b, G.Heusser^a, J.Hellmig^a, M. Hirsch^a, C. Hoffmann^a, H.V. Klapdor-Kleingrothaus^{*,a},
I.Kondratenko^b, D.Kotel'nikov^b, V.I.Lebedev^b, B.Maier^a, A.Müller^c, F.Petry^a, E.Scheer^a,
H.Strecker^a, M.Völlinger^a, K.Zuber^d

^aMax-Planck-Institut für Kernphysik, P.O.BOX 103980, 69029 Heidelberg, Germany

^bRussian Scientific Center-Kurchatov Institute, 123182 Moscow, Russia

^cINFN Gran Sasso, 67010 Assergi, Italy

^dInstitut für Hochenergiephysik, Schröderstr.90, 69120 Heidelberg, Germany

presented by K. ZUBER

Abstract

Recent results of the Heidelberg-Moscow double beta decay experiment are presented. After 8.6 kg·a of measuring time no signal is seen for the neutrinoless decay mode. A half-life limit of $T_{1/2}^{0\nu\beta\beta} > 5.1 \cdot 10^{24}$ a is deduced which converts into a neutrino mass limit of $\langle m_{\nu_e} \rangle < 0.68$ eV (90% CL). The experiment thus is the first one penetrating into the sub-eV range for the neutrino mass. For the 2ν mode a half life of $T_{1/2}^{2\nu\beta\beta} = (1.53 \pm 0.04_{\text{stat}} \pm 0.13_{\text{sys}}) \cdot 10^{24}$ a is derived. More than 10000 2ν double beta events are observed. This is the first high statistics observation of this nuclear decay mode. Limits on more exotic decay modes are also presented. Concerning dark matter the experiment now gives the sharpest limits for the observation of WIMPs.

1. INTRODUCTION

Neutrino physics has entered an era of new actuality. Several possible indications of physics beyond the Standard Model (SM) of particle physics are at present discussed: The lack of solar neutrinos, the atmospheric ν_μ deficit and mixed dark matter models could all be explained by non-vanishing ν -masses. Recent extended SO(10) scenarios with an S_4 horizontal symmetry which could explain these observations would predict degenerate ν -masses within 1-2 eV [1]. At present $\beta\beta$ -decay research may play a decisive role, since with

second generation $\beta\beta$ experiments like the Heidelberg-Moscow experiment using large amounts of enriched $\beta\beta$ -emitter material such a prediction can be tested in very near future.

The following decay modes for double beta decay of a nucleus $^Z_A X$ are usually considered:

$$2\nu\beta\beta : \quad ^Z_A X \rightarrow ^{Z+2}_A X + 2e^- + 2\bar{\nu}_e \quad (1)$$

$$0\nu\beta\beta : \quad ^Z_A X \rightarrow ^{Z+2}_A X + 2e^- \quad (2)$$

$$0\nu\chi\beta\beta : \quad ^Z_A X \rightarrow ^{Z+2}_A X + 2e^- + \chi \quad (3)$$

$$0\nu 2\chi\beta\beta : \quad ^Z_A X \rightarrow ^{Z+2}_A X + 2e^- + 2\chi \quad (4)$$

A positive evidence of processes (2)-(4) would require massive Majorana neutrinos, in addition a contribution

* Spokesmen of the collaboration

from right-handed weak currents is possible. The quantity which can be extracted out of process (2) is the effective Majorana mass given by

$$\langle m_{\nu_e} \rangle = \sum_i |U_{ei}^2 m_i| \quad (5)$$

where m_i are the mass eigenstates. Though there exists the possibility for destructive interference, in most grand unified models the effective mass seems to be equal to the electron neutrino mass eigenstate [2].

Process (3) would reflect breaking of a global (B-L) symmetry. The Majoron χ would be the Goldstone-boson associated with this symmetry breaking. Process (4) with the emission of two majorons seems possible in the case of supersymmetric models [3], and the same spectral shape for the sum energy of the electrons is expected if the majoron carries a leptonic charge [4]. The isotope under study at the Heidelberg-Moscow experiment [5] is ^{76}Ge in form of low-level semiconductor detectors. Further topics which can be investigated with such detectors are, for example, charge non-conservation [6] and dark matter [7].

2. THE EXPERIMENT

The Q -value of the decay to ^{76}Se is $Q = 2038.56$ keV. The experiment has 16.9 kg of Ge enriched to 86% in ^{76}Ge at hand (natural abundance of ^{76}Ge is 7.8 %). At present five enriched HP-detectors have been built from this material with a total mass of 11 kg of Ge. Table 1 gives some characteristics of the three detectors at present in regular operation at the Gran Sasso Laboratory. The given background B is determined between 2000-2080 keV. The next two detectors will be installed in fall 1994. The whole experiment is located underground in the Gran Sasso Laboratory in Italy (shielding depth is about 3500 m.w.e.). The detectors are built

Detector	Active mass [kg]	Enrichment [%]	Background [C/(kg·keV·a)]
Enr.1	0.920	86	0.16
Enr.2	2.758	86	0.21
Enr.3	2.324	88	0.22

Table 1. A comparison of the different detectors used in the experiment

in a 10 cm shield containing ultrapure LC2-lead surrounded by another 20 cm box of very pure Boliden lead. The whole setup is plugged in an air-free box which is flushed with high purity nitrogen. All parts of the detector cryostats are well selected and cleaned and have specific activities of less than $10 \mu\text{Bq/kg}$. To check the stability of the experiment a calibration is done every

Figure 1. The energy range between 2000-2080 keV of the spectrum seen after 8.6 kg·a of measurement. The dotted curve shows the peak excluded with 90% CL.

Figure 2. Remaining spectrum of enriched detector 2 after subtraction of all known background components (solid histogram, measuring time 19.2 mol·a). It is well fitted by a 2ν -spectrum (dashed line). Also shown is the measured spectrum stripped of all identified peaks (dotted histogram).

week with a ^{228}Th source.

3. RESULTS

Neutrinoless double beta decay

Fig.1 shows the region of interest of the $\beta\beta$ -spectrum after a measuring time of 8.6 kg·a. No line is seen so far. For a signal hidden in the background a half-life limit for this decay mode of $T_{1/2}^{0\nu} > 5.1(8.6) \cdot 10^{24}$ a (90% and 68% CL) can be derived. Using the matrix-elements given by [8] this can be converted into a limit for the neutrino mass of

$$\langle m_{\nu_e} \rangle < 0.68(0.52) \text{ eV} \quad 90\%(68\%) \text{ CL} \quad (6)$$

This is the most stringent limit for a neutrino mass coming out of double beta decay so far.

A transition to the first excited state would be dominated by right-handed currents and not by the mass-term like the ground state transition. For this transition we obtain a half-life limit to the first excited state of $T_{1/2}^{(0^+ \rightarrow 2^+)}(^{76}\text{Ge}) > 6.5 \cdot 10^{23}$ a (90% CL).

2ν double beta decay

In contrast to the neutrinoless decay the 2ν -mode has been observed in several isotopes, and there is also some evidence for an observation in ^{76}Ge . This decay mode can give information on nuclear structure. With the help of a Monte-Carlo simulation we subtracted all background components which can be identified, localized and quantitatively determined. The remaining counts are well fitted by a $2\nu\beta\beta$ -spectrum (Fig. 2). Using a maximum likelihood fit, we derive a half life of $T_{1/2}^{2\nu} = (1.53 \pm 0.04_{\text{stat}} \pm 0.13_{\text{sys}}) \cdot 10^{21}$ a after a measuring time of 49.6 mol·a (see also [9]). The number of $\beta\beta$ events is about 13500 and this decay is the main component of the measured spectrum in the region 500-1500 keV.

Majoron-accompanied double beta decay

For Majoron-accompanied $0\nu\beta\beta$ decay we look by further subtraction of the $2\nu\beta\beta$ spectrum from the

Figure 3. Exclusion limits for WIMPs with Ge detectors (dashed area). The increase of the sensitivity expected for some future [11] with enriched ^{73}Ge cryo detectors is also indicated (curves 99 and 99.9%). Also shown are theoretical expectations for spin-dependently interacting WIMPs from different GUT models.

Figure 4. Present situation and perspectives of the most promising $\beta\beta$ experiments. Only for the isotopes shown $0\nu\beta\beta$ half life limits $> 10^{21}$ a have been obtained. The thick solid lines correspond to the present, 1993, situation, open bars and dashed lines to 'safe' and 'less safe' expectations for 1999 .

'remaining spectrum' of Fig. 2. We obtain a half-life limit for this decay mode of $T_{1/2}^{0\nu\chi} > 7.81 \cdot 10^{21}$ a (90 % CL). This half-life limit can be converted into a limit for the neutrino-majoron-coupling constant $\langle g_{\nu\chi} \rangle$, since

$$T_{\frac{1}{2}}^{-1}[\text{a}^{-1}] = |M_{\text{GT}} - M_{\text{F}}|^2 F^{0\nu\chi} |\langle g_{\nu\chi} \rangle|^2 \quad (7)$$

where

$$\langle g_{\nu\chi} \rangle = \sum_{ij} g_{\nu\chi} U_{ei} U_{ej} \quad (8)$$

resulting in $\langle g_{\nu\chi} \rangle < 2.4 \cdot 10^{-4}$. For details see [10]. The set of matrix elements given by [8] is used. An investigation of the $0\nu 2\chi\beta\beta$ -mode results in a half-life limit of $T_{1/2}^{0\nu\chi\chi} > 7.05(8.54) \cdot 10^{21}$ a (90 % CL and 68 %CL).

Dark matter

Looking at dark matter interactions via coherent scattering of WIMPs off Ge nuclei requires sensitivity at low energies. By complementarily using an enriched ^{73}Ge detector, one would have a clear separation between spin-independent and spin-dependent interaction (^{73}Ge is the only germanium isotope with spin). After 165 kg·d of measurement with a lower threshold of 10 keV we obtained the exclusion plot shown in Fig. 3. The background is lower by a factor of 5-7 than that of other Ge-experiments dedicated to dark matter search, therefore improving the limits for heavier WIMPs [7, 12].

4. CONCLUSION and FUTURE

Using about 10 kg of enriched material, the final goal of the Heidelberg-Moscow experiment will be a half-life limit of $T_{1/2}^{0\nu} > 10^{25}$ a corresponding to neutrino masses down to about 0.1 eV. Thus it will be possible to check beyond the mentioned $\text{SO}(10) \times \text{S}_4$ model some other left-right symmetric see-saw models. The sensitivity of running and planned experiments till the end of the century is shown in Fig. 4. It is obvious that for making a significant step beyond the limit of about 0.1 eV in

reach of the present most sensitive experiments, *very* large increases of set-ups are necessary (see e.g. [13, 14])

Acknowledgements:

The experiment is supported by the Bundesministerium für Forschung und Technologie, the State Committee of Atomic Energy of Russia and the INFN. We thank the INFN Gran Sasso and especially Profs. P. Monacelli, E. Bellotti and N. Cabibbo for their generous support.

References

- [1] Lee, D.G., Mohapatra, R.N., preprint UMD-PP-94-95 (Feb. 1994)
- [2] Langacker, P. in *Neutrinos* ed. H.V. Klapdor, Springer 1988
- [3] Mohapatra, R.N., Takasugi, E., Phys. Lett. B **211**, 192 (1988)
- [4] Burgess, C.P., Cline, J.M., Phys. Lett. B **298**, 141 (1993)
- [5] Klapdor-Kleingrothaus, H.V., Proceedings of "Neutrinos in Cosmology, Astro-, Particle- and Nuclear Physics, Prog. Part. Nucl. Phys. **32** (1994) 261
- [6] Balysh, A. *et al.*, Phys. Lett. B **298**, 278 (1993)
- [7] Beck, M. *et al.*, Phys. Lett. B, in press (1994)
- [8] Staudt, A., Muto, K., Klapdor-Kleingrothaus, H.V., Europhys. Lett. **13**, 31 (1990);
Hirsch, M. *et al.*, Z. Phys. A **345**, 163 (1994);
Hirsch, M. *et al.*, Phys. Rep. **242**, 403 (1994)
- [9] Balysh, A. *et al.*, Phys. Lett B **322**, 176 (1994)
- [10] Beck, M. *et al.*, Phys. Rev. Lett. **70**, 2853 (1993)
- [11] Sadoulet, B., Nucl. Phys B (Proc. Suppl.) **35** (1994) 117
- [12] Beck, M., Nucl. Phys. B (Proc. Suppl.) **35**, 150
- [13] Raghavan, R.S., Phys. Rev. Lett. **10** (1994) 1411
- [14] Moe, M.K., Phys. Rev. C **44** (1991) R931